

Kalipanova I. Some cases from practice.

1. The boy of 8 years. Diagnosis: coxarthrosis of the right hip joint from 5 years. He was repeatedly treated in the hospital of Yaroslavl, received out-patient treatment in Rybinsk. There was no effect. When the boy turned to us, he get about on crutches, the right leg shorter than the left by 3cm, he had atrophy of shank, hip muscles, and complete atrophy of the right gluteus muscle. After 9 procedures of SCENAR-therapy in dosed mode, the influence was made on the both extremities, buttocks, directly on the joint and on three tracks), the legs became of the same length, atrophy of shank, hip, buttocks muscles disappeared. During the 12th and the 13th procedures, according SCENAR- expertise – there was a norm.

The second course was conducted in a month. The treatment after that was stopped because, objectively, no pathology was revealed. At control examination in Yaroslavl the pathology was not found but... diagnostic decision: the child must walk on crutches for a period of half of a year.

2. The girl of 9 years. Optic atrophy (left eye). Vis OS/OD=-0,001 /1.0. The girl could only discern by her left eye light and darkness. She was examined in St.-Petersburg, Moscow. Prognosis: hopeless, can't be cured. In 20 procedures of SCENAR-therapy the girl could read by her left eye a printed text within the distance of 10 cm from the eye, on the distance of 3 m she saw the first line in ophthalmologic table. The treatment continues.

3. 2 men turned to us because of myopia: one had – Vis OD / OS= - 6.0 / - 6.0, the other had: Vis OD / OS= - 3, 5 / -3, 5. In a month (3-4 procedures a week) of SCENAR-therapy, the first had Vis OD / OS= - 4, 0/-4, 0 and Vis OD / OS= - 1, 0 / -1, 0. In 15 more procedures - Vis OD / OS=-2.0 / -2, 0 and Vis OD / OS = 1, 0 / 1, 0. The treatment was finished before the departure to the conference. The influence was carried out locally and on cervicocollar zone.

4. The boy of 3 years. Diagnosis - infantile cerebral paralysis, spastic tetraparesis, mental retardation. During examination: the child couldn't hold his head, could only turn it to the right and to the left, cried, if he was lying on the stomach, he could only lie on the back. The backbone is flexible as if it is elastic; because of that the child couldn't stand. Spastic paresis of legs and arms. The child often repeats the words, which he hears, but forgets them at once. The influence was conducted on the extremities, body, in dosed mode. In 1,5 months of SCENAR-therapy the child walks himself with somebody, helping him, hold his head, creeping, but more than that, he likes to walk, he also likes to speak, speaks about the events, which take place at home. Arms paresis disappeared completely; there is only an insignificant paresis of the legs. It is marked that children with this disease undergo treatment very well, and the sooner the treatment started, the better it is.

The boy of 7 years. Chernobyl consequences. Examination showed: physical development is on the level of 4-5 years (when he was 5 years old, the growth stopped), says simple words with difficulty, but can't understand their meaning, doesn't know the meaning of the words "yesterday", "today", "tomorrow". Walks badly, can't run. The treatment was carried out with the help of SCENAR-532 and SCENAR-035-3 in individually-dosed mode. Taking into account the reason of the disease, the treatment was carried out from head to the toes of the feet. After the influence with the help of SCENAR-035-3 in addition therapy with the help of the healing blanket OLM-1 was made. The first course was 15 procedures with interval for the week-end. After that course of treatment the boy started running, grew up for 1 cm, the speech is distinct, he began to tell about the events, observing chronology.

In a month there was a repeated course of treatment, consisting of 10 procedures, the boy became 1 cm higher, the speech is more distinct, and he became communicative. It was decided to continue the treatment in 1-2 months.